DRAFT VERSION FEBRUARY 2, 2008
Preprint typeset using LaTeX style emulateapj v. 6/22/04

arXiv:astro-ph/0409766v1 30 Sep 2004

RADIAL VELOCITY DETECTABILITY OF LOW MASS EXTRASOLAR PLANETS IN CLOSE ORBITS

RAMAN NARAYAN, ANDREW CUMMING[1,2], AND D. N. C. LIN
Department of Astronomy and Astrophysics, University of California, Santa Cruz, CA 95064

Draft version February 2, 2008

ABSTRACT

Detection of Jupiter mass companions to nearby solar type stars with precise radial velocity measurements is now routine, and Doppler surveys are moving towards lower velocity amplitudes. The detection of several Neptune-mass planets with orbital periods less than a week has been reported. The drive toward the search for close-in Earth-mass planets is on the agenda. Successful detection or meaningful upper limits will place important constraints on the process of planet formation. In this paper, we quantify the statistics of detection of low-mass planets in-close orbits, showing how the detection threshold depends on the number and timing of the observations. In particular, we consider the case of a low-mass planet close to but not on the 2:1 mean motion resonance with a hot jupiter. This scenario is a likely product of the core-accretion hypothesis for planet formation coupled with migration of jupiters in the protoplanetary disk. It is also advantageous for detection because the orbital period is well-constrained. We show that the minimum detectable mass is $\approx 4\, M_\oplus\, (N/20)^{-1/2}(\sigma/\mathrm{m\ s^{-1}})(P/\mathrm{d})^{1/3}(M_\star/M_\odot)^{2/3}$ for $N \geq 20$, where N is the number of observations, P the orbital period, σ the quadrature sum of Doppler velocity measurement errors and stellar jitter, and $M_\star$ the stellar mass. Detection of few Earth mass rocky cores will require ~ 1 m s^{-1} velocity precision, and most important, a better understanding of stellar radial velocity "jitter".

Subject headings: planetary systems

1. INTRODUCTION

Over 100 Jupiter-mass planets have been discovered around nearby stars with precise radial velocity measurements (see Marcy et al. 2003 for a review). As time goes by, the range of periods and amplitudes accessible to these surveys increases, moving to longer periods and lower amplitudes (e.g. Carter et al. 2003; Fischer et al. 2003; Jones et al. 2003). Recently, the detection of Neptune-mass planets with periods less than ten days has been reported (Butler et al. 2004; McArthur et al. 2004; Santos et al. 2004). In this paper, we quantify the detectability of low mass planets in close orbits, and discuss the prospects for detecting earth mass objects. We first discuss possible origins of such planets and the importance of detecting them.

1.1. *The origin and importance of low mass rocky cores in close orbits*

In the conventional planet formation scenario (Safronov 1969; Pollack et al. 1996), heavy elements coagulate into terrestrial-planet-like cores prior to the formation of Jupiter-mass planets by subsequent accretion of gas. The critical mass which segregates these two populations is $M_c \sim$ a few $M_\oplus$, determined by the requirement of sufficiently high cooling efficiency in the gas envelope (Stevenson 1982; Bodenheimer & Pollack 1986; Ikoma et al. 2000). In a minimum mass nebula, both type of planets are thought to form at distances from their host stars comparable to those of planets in the solar system.

As a consequence of their tidal interaction, gas giant planets may undergo orbital migration (Goldreich & Tremaine 1980; Lin & Papaloizou 1986) which is finally halted either by their interaction with their host stars or through the local or global depletion of disks (Lin et al. 1996; Trilling et al. 2002; Armitage et al. 2002; Ida & Lin 2004). In this process, many gas giants are expected to be disrupted, transferring their mass onto the host star (Trilling et al. 2002), perhaps driven by tidal inflation (Gu, Lin, & Bodenheimer 2003). Ida & Lin (2004) estimate that 90–95% of planets that migrate to $a \lesssim 0.05$ AU must perish. Recent models of evaporation of material from hot jupiters (Lammer et al. 2003; Baraffe et al. 2004; Lecavalier des Etangs et al. 2004) find that gas giants may undergo significant mass loss, and perhaps be completely evaporated, during their lifetimes, although the efficiency and rates of evaporation remain uncertain (Yelle 2004). This inference raises the possibility of a class of remnant "hot Neptunes" or rocky cores in close orbits[3].

An alternative way to produce low-mass planets in close orbits is by resonant capture during migration of a hot jupiter. Along their migration paths, gas giant planets capture other planets onto their mean motion resonances as in the case of GJ 876 (Lee & Peale 2002; Nelson & Papaloizou 2002; Kley et al. 2004). Mandell & Sigurdsson (2003) also consider the survivability of earth mass objects in the wake of Jupiter migration. A natural implication of the core-accretion followed by migration scenario for the origin of hot jupiters is that they may harbor earth-mass planets ("hot earths") orbiting close to their mean motion resonances (Aarseth & Lin 2004, in preparation). In the gravitational instability scenario, however, both ice giants and terrestrial planets are assumed to emerge long after the formation of gas giant planets (Boss, Wetherill, & Haghighipour 2002).

Due to the effect of tidal circularization, the semimajor axis of the hot earth is expected to be slightly interior to the mean motion resonance, and its survival is ensured by the relativistic precession induced by the gravitational potential of the host star (Novak et al. 2003, Mardling & Lin 2004). After halting its inward migration, a hot jupiter may move outwards due to either its tidally induced mass loss (Trilling et al. 2002;

---

[1] Hubble Fellow

[2] Current address: Department of Physics, McGill University, 3600 rue University, Montreal, QC, Canada H3A 2T8.

[3] In this paper, we use the adjective "hot" to describe a closely-orbiting planet (orbital period $\lesssim 10$ days).

Gu et al. 2003), or tidal interaction with a rapidly spinning host star (Stassun et al. 1999; Dobbs-Dixon et al. 2004). In both cases, the periods of the close-in earth and hot jupiter may not be close to commensurability.

The presence of earth-mass planets close to the external mean motion resonance of a hot jupiter is also possible. In the wake of a migrating gas giant, the prototerrestrial planets in the disk external to its orbit may also undergo type I inward migration (Ward 1986) as a consequence of their own tidal interaction with the disk gas (Goldreich & Tremaine 1980). By inducing the formation of a gap near its orbit, a migrating jupiter provides a tidal barrier which prevents any migrating terrestrial planets from passing it. But this possibility is uncertain since type I migration is strongly affected by the poorly known turbulence near the coorbital region (Koller et al. 2003; Nelson & Papaloizou 2004). If this process is efficient, it could lead to the formation of an isolated hot earth (Ward 1997; Bodenheimer et al. 2000). Even if type I migration is inefficient, the presence of a hot jupiter can induce planetesimals to accumulate and form a hot earth just beyond the outer edge of the gap (Bryden et al. 2000). In the presence of additional long-period gas giant planets, a sweeping resonance may also lead to the inward migration and accumulation of terrestrial planets just exterior to the orbits of hot Jupiters (Lin, Nagasawa, & Thommes 2004).

Based on the core-accretion scenario for planet formation, it is already tempting to infer the existence of terrestrial planets in extrasolar systems from the detection of jupiter-mass gas giants (Ida & Lin 2004). However, the actual detection of a terrestrial mass object would be the first confirmation of rocky cores in an extrasolar planetary system (see however Kuchner 2003). The association of a hot earth with the interior mean motion resonance of a hot jupiter is particularly interesting because it would indicate that cores are formed prior to the emergence and migration of gas giant planets, providing strong support for the conventional core-accretion scenario for planet formation, and for the ubiquity of terrestrial planets in extrasolar planetary systems. The presence of a hot earth exterior to the orbit of a hot jupiter would support the notion of a tidal barrier which has the potential to enhance the formation probability of multiple planet systems. Finally, the discovery of any isolated hot earth would support the concept of type I migration.

1.2. *Outline of this paper*

In this paper, we discuss the detectability of such low mass companions in radial velocity surveys. In particular, we concentrate on the scenario in which a terrestrial planet was captured into the mean motion resonance of a migrating jupiter-mass planet, resulting in the formation of a "hot earth" which is near the interior mean motion resonances of a "hot jupiter". With the limitations of radial velocity surveys, the detection of a hot earth external to the orbit of a hot jupiter is much more difficult. We also calculate the detection thresholds for isolated low mass planets.

Being offset from the mean motion resonance of any nearby hot Jupiters, the expected contribution to the radial velocity amplitude K from a closely orbiting several Earth-mass object is

$$K = 6.4\ \mathrm{m/s}\ \left(\frac{M_P \sin i}{10\,M_\oplus}\right)\left(\frac{P}{\mathrm{days}}\right)^{-1/3}\left(\frac{M_\star}{M_\odot}\right)^{-2/3}, \qquad (1)$$

where P is the orbital period, M_P is the mass of the planet, and $M_\star$ is the mass of the star. This is comparable to the expected velocity variability from measurement errors, and intrinsic stellar "jitter" (Saar, Butler, & Marcy 1998). The orbital frequency is well-predicted for a hot earth close to a mean motion resonance with a hot jupiter, and so we expect the detectability to be somewhat better than a blind frequency search. Here we calculate the detectability and discuss the number of observations, and observing strategy needed for detection.



FIG. 1.— Simulated velocity residuals (top panel) after subtraction of a 0.5 M_J mass planet with $P = 3.1$ days, $K = 70$ m/s. We simulate one observation per night for 30 nights, with measurement error of 1 m s^{-1}. Periodogram (lower panel) evaluated for periods from 1 to 4 days before (dotted line) and after (solid line) subtraction. The strong feature present at ≈ 1.5 days before subtraction is due to aliasing of the 3.1 day signal. Subtracting reveals the presence of a 3 $M_\oplus$ companion at $P = 1.4$ days. The dashed line shows the detection threshold.

2. DETECTABILITY OF CLOSE COMPANIONS

2.1. *Search Technique*

We simulate a set of observational data for a system of planets which are on circular orbits. We search for periodicities using the Lomb-Scargle periodogram (Lomb 1976; Scargle 1982). For a set of N observation times $\{t_j\}$, velocities $\{v_j\}$, and measurement errors $\{\sigma_j\}$, and a trial orbital frequency $\omega = 2\pi/P$, we fit the function

$$f_j = A\cos\omega t_j + B\sin\omega t_j + C \qquad (2)$$

to the velocity data by minimising $\chi^2_\nu = (1/\nu)\sum_j (v_j - f_j)^2/\sigma_j^2$. The number of degrees of freedom is $\nu = N-3$ since there are 3 parameters in the model. We measure the goodness of fit as a function of trial frequency ω using the periodogram power z, defined as

$$z(\omega) = \frac{\Delta\chi^2/2}{\chi^2_\nu}, \qquad (3)$$

where $\Delta\chi^2 = (N-1)\chi^2_{N-1} - \nu\chi^2_\nu$, and χ^2_{N-1} is the reduced χ^2 of a fit of a constant to the data, $(N-1)\chi^2_{N-1} = \sum_j (v_j - \langle v\rangle)^2/\sigma_j^2$. Here, we extend the original Lomb-Scargle periodogram by allowing the mean to float at each frequency (Walker et al. 1995; Nelson & Angel 1998; Cumming, Marcy, & Butler 1999), rather than subtracting the mean of the data $\langle v\rangle$ prior to the fit. The periodogram power measures the improvement of

χ^2 when the sinusoid is included in the fit, in a similar way to a classical F-test (see Cumming 2004 for a recent discussion).

A typical Doppler measurement error is 3–5 m/s, although the precision of Doppler surveys continues to improve towards ~ 1 m/s (Mayor et al. 2003; Butler et al. 2004). However, at these high precisions, stellar "jitter" becomes the limiting factor. Jitter is observed at the few m/s level, depending on stellar properties such as age, rotation, and level of magnetic activity (Saar et al. 1998; Santos et al. 2000). For simplicity, we add Gaussian noise with amplitude $\sigma = 1$ m/s throughout this paper, which represents both Doppler errors and stellar jitter. The mass which can be detected scales proportional to σ, so that our results can be rescaled to the appropriate value of σ. For example, a 5 $M_\oplus$ planet with $\sigma = 5$ m/s has the same detectability as a 1 $M_\oplus$ planet with $\sigma = 1$ m/s. We have checked that this scaling applies for the two planet case, when the mass of the hot jupiter is much greater than the hot earth.

Our search procedure is to find the best-fit orbit for the hot jupiter, subtract it from the data, and search the residuals for periodicities. This is a much simpler approach than simultaneous fitting of both sinusoids, and is valid as long as the parameters from the two fits are uncorrelated. We have checked that this is the case, and find that only for sampling rates much greater than 1 day^{-1} is there some improvement by simultaneous fitting. An alternative possibility is that the orbital solution for the hot jupiter is already known from previous measurements, in which case this information can be used to subtract the hot jupiter's signal. Therefore, we also consider the cases where the full orbital solution or only the orbital period is known in advance.

If the hot jupiter orbital period is known in advance, we find the best fit amplitude and phase at that period. If no information is available, we search for the hot jupiter by evaluating $z(\omega)$ for periods between 2 and 4 days on a grid with frequency spacing $1/4T$, where T is the duration of the observations. To adequately subtract the hot jupiter's signal, it is important to accurately determine its orbital frequency. Therefore the frequency with the largest periodogram power is used as the starting value for a more accurate search for the maximum of $z(\omega)$. Once the hot jupiter's orbit has been subtracted, we calculate $z(\omega)$ for the residuals, and find the maximum periodogram power $z_{\rm max}$ near the 2:1 resonance. We do not include a constant term in the fit to the residuals, giving $\nu = N-2$. Figure 1 shows an example of the velocity residuals and the periodograms before and after subtraction. The peak at ≈ 1.5 days in the initial periodogram is due to aliasing of the hot jupiter's frequency. After subtraction, the signal due to a 3 $M_\oplus$ planet at 1.4 days shows clearly as a peak in z.

The importance of accurately determining the hot jupiter's orbital period is illustrated by Figure 2, which shows two examples of the velocity residuals after subtraction of the hot jupiter signal. In the first case (upper panels), we take the hot jupiter period returned by the periodogram (evaluated with a frequency spacing $1/4T$); in the second case (lower panels), we find a more accurate period estimate by searching for the peak in $z(\omega)$. In the first case, the velocity residuals are dominated by the inadequately-subtracted hot jupiter signal rather than the hot earth. How accurately must the hot jupiter period be known? If the orbital frequency is known to an accuracy $\delta\omega$, the amplitude of the residual part of the hot jupiter's signal is $\Delta v \approx A_1\delta\omega T$, where A_1 is the hot jupiter amplitude. The accuracy to which the hot jupiter frequency may be determined is $\delta\omega \approx (2\pi/T)(\sigma/\sqrt{N}A_1)$ (e.g. Bretthorst 1988), so that

$$\Delta v \approx \frac{2\pi\sigma}{\sqrt{N}}, \tag{4}$$

independent of T and A_1. For large N, $\Delta v \ll \sigma$, so that the residual part of the hot jupiter signal has no effect on detectability. However, for $N \lesssim (2\pi)^2$, the residuals from the subtraction contribute a significant additional source of velocity variability. Even when the hot jupiter period is known in advance, there is an uncertainty in the fitted amplitude A_1 of $\approx \sigma/\sqrt{N}$, and phase of $\approx (2/N)^{1/2}(\sigma/K)$, giving an additional velocity scatter for low N. Therefore for small N we expect detection of a hot earth to be more difficult when the full orbital solution for the hot jupiter is not specified in advance.

2.2. *Calculation of Detection Threshold and Detection Probabilities*

The significance of the maximum observed periodogram power $z_{\rm max}$ depends on how often an equally good or better fit would occur purely due to a noise fluctuation. We determine this with Monte Carlo simulations. We generate data sets with a hot jupiter plus noise, i.e. without a hot earth, and search for a second companion. The 99% detection threshold z_d is determined as the value of z which is exceeded in only 1% of trials, or alternatively for which there is a 1% false alarm probability F. The detection threshold is indicated in Figure 1 by the dashed line.

We adopt a similar Monte Carlo approach to determine the detection probability. We generate a large number of data sets for each choice of M_p and N, and calculate the fraction of trials for which the hot earth is detected. A random distribution of inclinations is included. Figure 3 shows the detection efficiency as a function of number of observations N for $M_P = 1$, 1.5, 2, and 3 $M_\oplus$. We simulate 1 observation per 8 hour night, and take observations for successive nights. In these simulations, we search for the hot earth orbital period between 1 and 1.8 days. Figure 4 shows the effect of changing this frequency range. If the frequency of the hot earth is specified in advance, the detectability is increased since the number of "independent frequencies" in the search is less, giving a smaller chance of a false alarm due to a noise fluctuation. This is the well-known "bandwidth penalty" (e.g. Vaughan et al. 1994).

In Figure 5, we show the number of observations needed to detect a given mass 50% of the time. The crosses are for data sets containing only a hot Earth, or equivalently, for the case when the hot jupiter orbit is completely known in advance. The triangles are for the case with a hot jupiter whose orbital period is known in advance, and the circles for the case with a hot jupiter, but with a full search for both orbital periods. For $N \approx 10$–20, planets with $M \gtrsim 4(\sigma/1\ {\rm m\ s^{-1}})\ M_\oplus$ can be detected. However, detection of a planet with $M \sim 1(\sigma/1\ {\rm m\ s^{-1}})\ M_\oplus$ requires $N \approx 200$. For $N \lesssim 20$, detection of the hot earth after fitting for the hot jupiter's signal is harder than detection of the hot earth alone or with the hot jupiter orbit fully specified in advance (compare the crosses and circles in Fig 5). This is likely due to the additional velocity scatter at low N from inadequate subtraction of the hot jupiter's signal ($\Delta v \gtrsim \sigma$ in eq. [4]).

2.3. *Analytic estimates*

In this section, we derive an analytic estimate for the detection threshold, following the approach of Cumming (2004),



FIG. 2.— The velocity residuals and periodogram after subtraction of the hot jupiter for two different estimates of the hot jupiter's orbital frequency. In the upper panel, we use the frequency estimate from the periodogram evaluated with frequency spacing $1/4T$. In this case, the poor subtraction leads to a large scatter in the residuals. In the lower panel, we determine an accurate estimate of the frequency by finding the peak of $z(\omega)$. The velocity residuals are then dominated by the 5 $M_\oplus$ companion.



FIG. 3.— Detection efficiency for hot earths with masses of 1, 1.5, 2, and 3 $M_\oplus$ and period 1.4 days, with σ = 1 m s^{-1}. We assume 1 randomly-timed observation on successive 8 hour nights. We use 10,000 trials to evaluate the detection probability. The hot Jupiter has a mass of 1.0 Jupiter mass, and a period of 3.0 days.



FIG. 4.— The effect of the range in orbital period searched on the detection probability. We take σ = 1ms^{-1} and M_P = 3 $M_\oplus$. Triangles represent orbital periods ranging from 1 – 4 days, squares 1.3 – 1.5 days, and circles 1.39 – 1.41 days. The dotted line is for 1 – 1.8 days, which is used elsewhere in this paper. As a larger frequency range is searched, N_i increases giving a larger detection threshold and lower detection probability.

who discusses the detectability of single planets with radial velocities. First, we determine z_d semi-analytically for Gaussian noise. The cumulative distribution of $z(\omega)$ for a single frequency is (e.g. Cumming et al. 1999)

$$\mathrm{Prob}(z > z_0) = \left(1 + \frac{2z_0}{\nu}\right)^{-\nu/2}, \tag{5}$$

or for large N,

$$\mathrm{Prob}(z > z_0) \approx \exp(-z_0),. \tag{6}$$

For a given false alarm probability F, the detection threshold is given by

$$F = 1 - [1 - \mathrm{Prob}(z > z_d)]^{N_i}, \tag{7}$$

where N_i is the number of "independent frequencies" searched. For $F \ll 1$,

$$F \approx N_i\ \mathrm{Prob}(z > z_d). \tag{8}$$

For unevenly-sampled data, N_i must be determined by Monte Carlo simulations. However, since the spacing of periodogram peaks is $1/T$, a rough estimate is $N_i \approx T\Delta f$, where Δf is the frequency range searched (Cumming 2004).



FIG. 5.— The number of observations needed to detect a planet 50% of the time as a function of the mass M_P scaled by the error σ. Observations are taken at a random time during each consecutive night. The crosses are for data sets containing only a hot Earth, the triangles for the case where the hot jupiter period is known in advance, and the circles for a full search for both orbital periods. The curves are the analytic results. The dotted curve occurs if the number of independent frequencies, N_i, is $N_i = T\Delta f$, where T is the duration and Δf is the frequency range used to search for the hot earth. For the dashed curve, $N_i = 10$.

We next make an analytic estimate of the detection threshold. In the presence of a signal with amplitude K, the average power is $z_s = \nu K^2/4\sigma^2$ (Groth 1975; Scargle 1982; Horne & Baliunas 1986, we have accounted for the different normalization). For a given N_i and F, we find the detection threshold z_d by inverting equation (7) using the analytic distribution (5). Then, setting $z_s = z_d$ gives the signal to noise ratio needed to detect the signal

$$\frac{K}{\sqrt{2}\sigma} = \left[\left(\frac{N_i}{F}\right)^{2/(N-3)} - 1\right]^{1/2} \tag{9}$$

$$\approx \left[\frac{2\ln(N_i/F)}{N}\right]^{1/2}, \tag{10}$$

(Cumming et al. 2003; Cumming 2004) where the approximation is for large N. As expected, the signal to noise ratio goes down as $1/\sqrt{N}$ for $N \gg 1$.

Combining equation (1) with equation (9) or (10) gives the velocity amplitude and mass which can be detected 50% of the time. Equation (10) ($N \gg 1$) gives

$$K_{50} = \frac{6\ \mathrm{m\ s^{-1}}}{\sqrt{N}} \left(\frac{\sigma}{\mathrm{m\ s^{-1}}}\right) \left(\frac{\ln[N_i/F]}{9.2}\right)^{1/2}, \tag{11}$$

or

$$M_{50} \approx \frac{10\ M_\oplus}{\sqrt{N}} \left(\frac{\sigma}{\mathrm{m\ s^{-1}}}\right) \left(\frac{P}{\mathrm{d}}\right)^{1/3} \left(\frac{\ln[N_i/F]}{9.2}\right)^{1/2} \left(\frac{M_\star}{M_\odot}\right)^{2/3}, \tag{12}$$

where we have also used the fact that the mean value of $\sin i$ is $\pi/4$, and we take $N_i = 100$ and $F = 0.01$.

Figure 5 compares the analytic results for arbitrary N (eq. [9]) with our numerical simulations. The dashed and dotted curves show the analytic models for two different estimates of N_i. The agreement is excellent for a single planet. For the two planet case, equation (12) applies for $N \gtrsim 20$, but underestimates M_{50} for $N \lesssim 20$. For such low N, we expect $\Delta\nu \gtrsim \sigma$ (eq. [4]), making detection of the hot earth more difficult than in the single planet case.

2.4. *Observing Strategy*

We now address to what extent the spacing of the observation times affects the detectability of the hot earth. First, we consider observations made on successive nights, but with different numbers of observations per night. Figure 6 shows the detection probability as a function of number of nights, and as a function of number of observations. For a fixed number of observations, increasing the sampling rate reduces the duration of the data, leading to less frequency resolution and therefore smaller N_i. We therefore expect better detectability for a faster sampling rate at a fixed number of observations. This is initially the case, as can be seen by comparing the curves for 1 per night and 5 per night in Figure 6. However, for very rapid sampling, a large number of observations must be made before a complete hot jupiter orbit is sampled, therefore allowing the orbital parameters to be determined accurately enough for a good subtraction. This is likely the reason for the decrease in detectability on going to more than 5 observations per night in Figure 6. Although observing more rapidly always leads to detection in a fewer number of nights (left panel of Fig. 6), the number of observations required eventiually becomes very large (right panel of Fig. 6). Therefore there is an optimum observing rate (roughly a few per night for this case, but it depends slightly on the hot earth mass).

Although observing on successive nights would be possible with a dedicated telescope, current radial velocity surveys are limited by telescope scheduling. Therefore, we have also simulated more realistic observation times for current surveys, by observing for three nights each month, with the seperation between observing runs randomly chosen between 20 and 40 days. We find that if the hot jupiter orbital parameters are not known in advance, the detectability can be affected by the observing scheme, because of aliasing of the hot jupiter frequency leading to subtraction of an incorrect orbit. However, in the practical case that the hot jupiter period is known from previous observations, the hot jupiter orbit can be adequately subtracted, and the effect of the observing strategy is small.

3. SUMMARY AND DISCUSSION

We have calculated the detectability of low mass planets in radial velocity surveys, and in particular a "hot earth" companion to a hot jupiter. Detection of such a companion would give important clues to the ordering of the planet formation process, and the ubiquity of terrestrial mass cores around solar type stars. The velocity amplitude and mass required for a 50% detection rate are given by equations (11) and (12), and shown in Figure 5. For $N \approx 20$, masses greater than $4\ M_\oplus\ (\sigma/\mathrm{m\ s^{-1}})(P/\mathrm{d})^{1/3}(M_\star/M_\odot)^{2/3}$ have 50% detectability or better.

Our results apply to a low mass planet that is near the mean motion resonance with a hot jupiter, or that is isolated. In Figure 7, we show the detection limits of equation (12) compared to the present distribution of known exoplanets[4]. The left panel is for $M_\star = 1\ M_\odot$, the right panel for $M_\star = 0.2\ M_\odot$. Lower mass stars have a greater velocity amplitude for a given planet mass, so that M dwarfs are the most promising to

---

[4] Taken from http://www.exoplanets.org/.



FIG. 6.— Detection probability for a 1 $M_\oplus$ planet with different numbers of observations made on consecutive nights. We show the detection probability against number of nights (left panel) and number of observations (right panel). A more rapid sampling rate leads to a detection in fewer nights, but at the cost of taking many observations.

TABLE 1
SUMMARY OF RADIAL VELOCITY DATA FOR HOT JUPITERS AND ESTIMATED UPPER LIMITS FOR HOT EARTHS

Star name	References	P_J[a] (d)	rms[b] (m s^{-1})	N	T (d)	N_i[c]	$M_\star$ ($M_\odot$)	K_{99}[d] (m s^{-1})	M_{99}[e] ($M_\oplus$)
HD 73256	1	2.55	15	40	80	4	1.05	22	58
HD 83443	2,3	2.99	3.8	36	1150	38	0.79	7.3	15
HD 46375	4	3.02	2.6	24	516	21	1.0	6.5	15
HD179949	5	3.09	10	23	735	30	1.24	27	74
HD 187123	6	3.10	7.6	20	250	10	1.0	20	48
τ Boo	7	3.31	16.5	58	4400	166	1.3	26	74
BD-103166	8	3.49	8.1	17	410	15	1.1	26	69
HD 75289	9	3.51	7.5	88	330	12	1.15	7.7	21
HD 76700	10	3.97	6.2	24	1243	39	1.0	16	42

[a] Orbital period of the hot jupiter.
[b] rms of the residuals to the orbital solution.
[c] Estimated number of independent frequencies for a search from 2–2.4 times the hot jupiter orbital frequency, $N_i \approx 0.1T/P_J$.
[d] Estimated 99% upper limit, using equation (9) with $K_{99} \approx 1.7K_{50}$.
[e] 99% upper limit on the mass of a hot earth near the 2:1 resonance with the hot jupiter.
[f] References.—(1) Udry et al. 2003; (2) Mayor et al. 2002; (3) Butler et al. 2002; (4) Marcy et al. 2000; (5) Tinney et al. 2001; (6) Butler et al. 1998; (7) Fischer et al. 2001; (8) Butler et al. 2000; (9) Udry et al. 2000; (10) Tinney et al. 2003

search for terrestrial mass bodies. Close orbits are particularly interesting for this case, since they lie within the habitable zone for M dwarfs (orbital periods of a few to tens of days; Kasting, Whitmire, & Reynolds 1993; Joshi, Haberle, & Reynolds 1997).

Recently, three Neptune-mass planets have been discovered in close orbits around GJ 436 (Butler et al. 2004; $M \sin i = 21\ M_\oplus$, $P = 2.644$d, $N = 42$, rms=5.3 m s^{-1}), ρ Cnc (McArthur et al. 2004; $M \sin i = 14.2\ M_\oplus$, $P = 2.808$d, $N = 119$, rms=5.4 m s^{-1}), and μ Ara (Santos et al. 2004; $M \sin i = 14\ M_\oplus$, $P = 9.5$d, $N = 24$, rms=0.9 m s^{-1}). These can be seen as the lowest mass planets in Figure 7. Only one of these stars, GJ 436, is an M dwarf. All of these detections lie above the detection limits in Figure 7.

The lowest mass planet detected prior to these recent announcements was HD 49674b (Butler et al. 2002) with $M \sin i = 0.12\ M_J$ (or $\approx 40\ M_\oplus$), $K = 13$ m s^{-1}, and $P = 5$ days. The velocity curve presented by Butler et al. (2002) has $N = 24$, and the residual rms is ≈ 5 m s^{-1}. With these values of N and σ, the thresholds we derive here imply that planets with half the mass of HD 49674b ($\approx$ 10–20 $M_\oplus$ depending on orbital period) should be detectable, consistent with the recent detections. Endl et al. (2003) report 22 velocity measurements of the M dwarf Proxima Cen with Doppler errors of 2.5 m s^{-1}. They calculate an upper limit of 4–6 $M_\oplus$ for planets in the habitable zone (estimated to be 4–14 days), in good agreement with Figure 7.

A summary of published data for hot jupiters with orbital periods $P_J < 4$ days is shown in Table 1. To estimate an upper limit on the velocity amplitude of a hot earth, we first use the duration and number of observations to estimate N_i for a search between orbital periods $P_J/2.4$ and $P_J/2$. We then write the 99% upper limit as $K_{99} = 1.7K_{50}$, where K_{50} is given by equation (9), but with the rms of the residuals to the best-



FIG. 7.— Summary of detection thresholds in the mass-period plane. The dashed lines show 50% detection thresholds for different N and σ, for $M_\star = 1\ M_\odot$ (left panel) and $M_\star = 0.2\ M_\odot$ (right panel). It is assumed that the duration of the observations is longer than the orbital period. The circles show currently detected planets ($M \sin i$) from exoplanets.org, and including the 3 recently announced Neptune-mass candidates. The dotted lines show velocity amplitudes of $K = 1, 3$, and 10 m/s ($\sin i = 1$), and the dashed line shows an approximate detection threshold for SIM (assuming 1μas sensitivity and a 10 pc distance; see Ford & Tremaine 2003).

fitting hot jupiter orbit substituted for σ. Since we adopt the full rms of the residuals as the noise level, these are conservative upper limits. The values of K_{99} range from 7–30 m s^{-1}, corresponding to $\approx$ 15–80 $M_\oplus$ for companions close to 2:1 resonance with the hot jupiter. HD 83443 and HD 46375 have upper limits of 15 $M_\oplus$ because of very precise measurements giving a low rms (4 and 3 m s^{-1} respectively).

These detections and upper limits imply that the current threshold for detecting planets in close orbits is $\gtrsim$ 10–20 $M_\oplus$, with planets close to this threshold just starting to be discovered. With more observations, and improvements in measurement precision, this threshold can be pushed lower. In Figure 7, we show the detectable mass for $N = 100$ and $\sigma = 3$ m s^{-1}, which reaches a few Earth masses for orbits $\sim$ 1 day for a solar mass star, or 1 $M_\oplus$ for a 0.2 $M_\odot$ M dwarf. However, a better understanding of stellar jitter will be crucial for convincing detections of small amplitude planets. Detection of extrasolar planets with radial velocities becomes more and more difficult as K approaches σ, because alternative explanations for the detected signal must be excluded (see the discussion in Cumming et al. 1999 for example). These include systematic errors in the measurements, as well as stellar jitter. Both may be periodic, with timescales associated with seasonal variations, or stellar properties such as rotation, convective motions, or the appearance and disappearance of magnetic features. Saar & Fischer (2000), Paulson et al. (2002), and Saar (2003) discuss attempts to correct for stellar jitter using simultaneous measurements of activity indicators or spectral line shapes, but such studies are just beginning. Very precise Doppler measurements with errors at the 1 m s^{-1} level will help to understand the stellar jitter and systematic errors.

Finally, we have assumed that both planets are in circular orbits in this paper. For orbital periods as short as a few days, the timescale for tidal circularization is expected to be much shorter than the age of the system. However, additional planets in the system, at longer orbital periods and so far undetected, may excite the eccentricity of the hot jupiter. Tidal damping of an eccentricity induced in this way has been suggested as the reason for the inflated radius of the transiting planet HD 209458b (Bodenheimer, Lin, & Mardling 2001). The required eccentricity is comparable to current measurement uncertainties, typically $\delta e \sim 0.03$. Fitting a circular orbit to a Keplerian orbit with eccentricity e gives a residual scatter $\approx eA_1$, where A_1 is the amplitude. For a hot jupiter with $A_1 \approx 100$ m s^{-1} the scatter in the residuals is comparable to the signal from a hot earth. However, since the Fourier components of a Keplerian orbit are at the orbital period and its harmonics, whereas the hot earth is expected to lie outside the 2:1 resonance, it should be possible to distinguish between these two possibilities, although simultaneous fitting of the hot jupiter and earth orbits may be required. Therefore we do not expect a significant reduction in sensitivity over the detection thresholds calculated in this paper. This issue is closely related to the important question of determining the uncertainty in orbital parameters (Ford 2003), particularly the eccentricity of hot jupiters, and deserves further study.

We acknowledge support from NSF through grant AST-9987417 and NASA through grant NAG5-13177. AC is supported by NASA through Hubble Fellowship grant HF-01138 awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS 5-26555.

REFERENCES

Armitage, P. J., Livio, M., Lubow, S. H., & Pringle, J. E. 2002, MNRAS, 334, 248

Baraffe, I., Selsis, F., Chabrier, G., Barman, T. S., Allard, F., Hauschildt, P. H., & Lammer, H. 2004, A&A, 419, L13

Bodenheimer, P. & Pollack, J. B. 1986, Icarus, 67, 391

Bodenheimer, P., Hubickyj, O. Lissauer, J. J. 2000, Icarus, 143, 2

Bodenheimer, P., Lin, D. N. C., & Mardling, R. A. 2001, ApJ, 548, 466

Boss, A. P., Wetherill, G. W., & Haghighipour, N. 2002, Icarus, 156, 291

Bretthorst, L. G. 1988, "Bayesian Spectrum Analysis and Parameter Estimation", Lecture Notes in Statistics vol. 48 (Springer-Verlag)

Bryden, G., Rzyczka, M., Lin, D. N. C., & Bodenheimer, P 2000, ApJ, 540, 1091

Butler, R. P., Marcy, G. W., Vogt, S. S., & Apps, K. 1998, PASP, 110, 1389

Butler, R. P., Vogt, S. S., Marcy, G. W., Fischer, D. A., Henry, G. W., & Apps, K. 2000, ApJ, 545, 504

Butler, R. P., et al. 2002, ApJ, 578, 565

Butler, R. P., Bedding, T. R., Kjeldsen, H., McCarthy, C., O'Toole, S. J., Tinney, C. G., Marcy, G. W., & Wright, J. T. 2004, ApJ, 600, L75

Butler, R. P., Vogt, S. S., Marcy, G. W., Fischer, D. A., Wright, J. T., Henry, G. W., Laughlin, G., & Lissauer, J. J. 2004, ApJ, submitted (astro-ph/0408587)

Carter, B. D., Butler, R. P., Tinney, C. G., Jones, H. R. A., Marcy, G. W., McCarthy, C., Fischer, D. A., & Penny, A. J. 2003, ApJ, 593, L43

Cumming, A. 2004, MNRAS, in press (astro-ph/0408470)

Cumming, A., Marcy, G. W., & Butler, R. P. 1999, ApJ, 526, 890

Cumming, A., Marcy, G. W., Butler, R. P., & Vogt, S. S. 2003, in ASP Conf. Ser. 294, Scientific Frontiers in Research on Extrasolar Planets, ed. D. Deming, & S. Seager (San Francisco: ASP), 27

Dobbs-Dixon, I., Lin, D.N.C., & Mardling, R.A. 2004, ApJ, in press

Endl, M., Kürster, M., Rouesnel, F., Els, S., Hatzes, A. P., & Cochran, W. D. 2003, in ASP Conf. Ser. 294, Scientific Frontiers in Research on Extrasolar Planets, ed. D. Deming, & S. Seager (San Francisco: ASP), 75

Fischer, D. A., Marcy, G. W., Butler, R. P., Vogt, S. S., Frink, S., & Apps, K. 2001, ApJ, 551, 1107

Fischer, D. A., Butler, R. P., Marcy, G. W., Vogt, S. S., & Henry, G. W. 2003, ApJ, 590, 1081

Ford, E. B. 2003, preprint (astro-ph/0305411)

Ford, E. B. & Tremaine, S. 2003, PASP, 115, 1171

Goldreich, P., & Tremaine, S. 1980, ApJ, 241, 425

Groth, E. J. 1975, ApJS, 29, 285

Gu, P., Lin, D. N. C., & Bodenheimer, P. H. 2003, ApJ, 588, 509

Horne, J. H. & Baliunas, S. L. 1986, ApJ, 302, 757

Ida, S. & Lin, D. N. C. 2004, ApJ, 604, 388

Ikoma, M., Nakazawa, K. & Emori, E. 2000, ApJ, 537, 1013

Jones, H. R. A., Butler, R. P., Tinney, C. G., Marcy, G. W., Penny, A. J., McCarthy, C., & Carter, B. D. 2003, MNRAS, 341, 948

Joshi, M. M., Haberle, R. M., Reynolds, R. T. 1997, Icarus, 129, 450

Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, Icarus, 101, 108

Kley, W., Peitz, J. & Bryden, G. 2004, A&A, 414, 735

Koller, J., Li, H., & Lin, D.N.C. 2003, ApJ, 596, 91

Kuchner, M. J. 2003, ApJ, 596, L105

Lammer, H., Selsis, F., Ribas, I., Guinan, E. F., Bauer, S. J., & Weiss, W. W. 2003, ApJ, 598, L121

Lecavalier des Etangs, A., Vidal-Madjar, A., McConnell, J. C., & Hebrard, G. 2004, A&A, 418, L1

Lee, M.H. & Peale, S. 2002, ApJ, 567 596

Lin, D. N. C., Bodenheimer, P. & Richardson, D. 1996, Nature, 380, 606

Lin, D. N. C., Nagasawa, M., & Thommes, E. 2004, Nature, submitted

Lin, D. N. C., & Papaloizou, J. C. B. 1986, ApJ, 309, 846

Lomb, N. R. 1976, Ap&SS, 39, 447

Mandell, A. M., & Sigurdsson, S. 2003, ApJ, 599, L111

Marcy, G. W., Butler, R. P., & Vogt, S. S. 2000, ApJ, 536, L43

Marcy, G. W., Butler, R. P., Fischer, D. A., & Vogt, S. S. 2003, in ASP Conf. Ser. 294, Scientific Frontiers in Research on Extrasolar Planets, ed. D. Deming, & S. Seager (San Francisco: ASP), 1

Mardling, R. & Lin, D.N.C. 2004, ApJ, submitted

Mayor, M., Naef, D., Pepe, F., Queloz, D., Santos, N. C., Udry, S., & Burnet, M. 2002, in ASP Conf. Ser., Planetary Systems in the Universe, ed. A. J. Penny, P. Artymowicz, A. M. Lagrange, & S. S. Russell (San Francisco: ASP), in press

Mayor, M., et al. 2003, The Messenger, 114, 20

McArthur, B. E. et al. 2004, ApJ, in press (astro-ph/0408585)

Nelson, A. F. & Angel, J. R. P. 1998, ApJ, 500, 940

Nelson, R. P., & Papaloizou, J. C. B. 2002, MNRAS, 333, 26

Nelson, R. P., & Papaloizou, J. C. B. 2004, MNRAS, 350, 849

Novak, G. Lai, D., & Lin, D.N.C. 2003, in ASP Conf. Ser. 294, Scientific Frontiers in Research on Extrasolar Planets, ed. D. Deming, & S. Seager (San Francisco: ASP), 177

Novak, G. Lai, D., & Lin, D.N.C. 2004, ApJ, submitted

Paulson, D. B., Saar, S. H., Cochran, W. D., & Hatzes, A. P. 2002, ApJ, 124, 572

Pollack, J. B., Hubickyj, O., Bodenheimer, P., Lissauer, J. J., Podolak, M., & Greenzweig, Y. 1996, Icarus, 124, 62

Saar, S. H. 2003, in ASP Conf. Ser. 294, Scientific Frontiers in Research on Extrasolar Planets, ed. D. Deming, & S. Seager (San Francisco: ASP), 65

Saar, S. H., Butler, R. P., & Marcy, G. W. 1998, ApJ, 498, L153

Saar, S. H. & Donahue, R. A. 1997, ApJ, 485, 319

Saar, S. H. & Fischer, D. 2000, ApJ, 534, L105

Safronov, V. 1969, Evolution of thr Protoplanetary Cloud and Formation of the Earth and Planets (Moscow: Nauka Press)

Santos, N. C., Mayor, M., Naef, D., Pepe, F., Queloz, D., Udry, S., & Blecha, A. 2000, A&A, 361, 265

Santos, N. C., et al. 2004, A&A, submitted (astro-ph/0408471)

Scargle, J. D. 1982, ApJ, 263, 835

Schwarzenberg-Czerny, A. 1998, Baltic Astron., 7, 43

Stevenson, D. J. 1982, P&SS, 30, 755

Stassun, K. G., Mathieu, R. D., Mazeh, T., & Vrba, F. J. 1999, AJ, 117, 2941

Tinney, C. G., Butler, R. P., Marcy, G. W., Jones, H. R. A., Penny, A. J., McCarthy, C., Carter, B. D., & Bond, J. 2003, ApJ, 587, 423

Tinney, C. G., Butler, R. P., Marcy, G. W., Jones, H. R. A., Penny, A. J., Vogt, S. S., Apps, K., & Henry, G. W. 2001, ApJ, 551, 507

Trilling, D. E., Lunine, J. I. & Benz, W. 2002, A&A, 394, 241

Udry, S., et al. 2000, A&A, 356, 590

Udry, S., et al. 2003, A&A, 407, 679

Vaughan, B. A., et al. 1994, ApJ, 435, 362

Walker, G. A. H., Walker, A. R., Irwin, A. W., Larson, A. M., Yang, S. L. S., & Richardson, D. C. 1995, Icarus, 116, 359

Ward, W.R. 1986, Icarus, 67, 164

Ward, W.R. 1997, ApJ, 482, L211

Yelle, R. V. 2004, Icarus, in press